

## Additional exercises for Book D

---

### Contents

Additional exercises for Unit D1	2
Solutions to additional exercises for Unit D1	4
Additional exercises for Unit D2	8
Solutions to additional exercises for Unit D2	10
Additional exercises for Unit D3	15
Solutions to additional exercises for Unit D3	16
Additional exercises for Unit D4	20
Solutions to additional exercises for Unit D4	22

# Additional exercises for Unit D1

## Section 1

### Additional Exercise D1

Arrange the following numbers in increasing order.

- (a)  $7/36$ ,  $3/20$ ,  $1/6$ ,  $7/45$ ,  $11/60$   
 (b)  $0.\overline{465}$ ,  $0.4\overline{65}$ ,  $0.46\overline{5}$ ,  $0.4655$ ,  $0.4656$

### Additional Exercise D2

Find the fractions whose decimal representations are as follows.

- (a)  $0.\overline{481}$       (b)  $0.48\overline{1}$

### Additional Exercise D3

Let  $x = 0.\overline{21}$  and  $y = 0.\overline{2}$ . Use long division to find  $x + y$  and  $xy$  in decimal form.

### Additional Exercise D4

Find a rational number  $x$  and an irrational number  $y$ , both in the interval  $(0.119, 0.12)$ .

## Section 2

### Additional Exercise D5

Solve the following inequalities.

- (a)  $\frac{x-1}{x^2+4} < \frac{x+1}{x^2-4}$       (b)  $\sqrt{4x-3} > x$   
 (c)  $|17 - 2x^4| \leq 15$

## Section 3

### Additional Exercise D6

Use the Triangle Inequality to prove that

$$|a| \leq 1 \implies |a-3| \geq 2.$$

### Additional Exercise D7

Prove that

$$(a^2 + b^2)(c^2 + d^2) \geq (ac + bd)^2, \text{ for } a, b, c, d \in \mathbb{R}.$$

### Additional Exercise D8

Prove the inequality

$$3^n \geq 2n^2 + 1, \text{ for } n \geq 1,$$

- (a) by using the Binomial Theorem, applied to  $(1+x)^n$  with  $x=2$   
 (b) by using mathematical induction.

### Additional Exercise D9

Apply Bernoulli's Inequality, first with  $x = 2/n$  and then with  $x = -2/(3n)$ , to prove that

$$1 + \frac{2}{3n-2} \leq 3^{1/n} \leq 1 + \frac{2}{n}, \text{ for } n \geq 1.$$

### Additional Exercise D10 Challenging

Use mathematical induction to prove that

$$\frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} \leq \sqrt{\frac{3/4}{2n+1}}, \text{ for } n \geq 1.$$

## Section 4

### Additional Exercise D11

In this exercise, take

$$E_1 = \{x \in \mathbb{Q} : 0 \leq x < 1\}$$

and

$$E_2 = \{(1 + 1/n)^2 : n = 1, 2, \dots\}.$$

- (a) Prove that each of the sets  $E_1$  and  $E_2$  is bounded above. Which of them has a maximum element?
- (b) Prove that each of the sets  $E_1$  and  $E_2$  is bounded below. Which of them has a minimum element?
- (c) Determine the least upper bound of each of the sets  $E_1$  and  $E_2$ .
- (d) Determine the greatest lower bound of each of the sets  $E_1$  and  $E_2$ .

# Solutions to additional exercises for Unit D1

## Solution to Additional Exercise D1

(a) Putting all the fractions over the common denominator of 180, we obtain

$$\frac{3}{20} < \frac{7}{45} < \frac{1}{6} < \frac{11}{60} < \frac{7}{36}.$$

(Alternatively, we can compare the decimal representations of these fractions.)

(b)  $0.\overline{465} < 0.4655 < 0.4\overline{65} < 0.4656 < 0.4\overline{65}$

## Solution to Additional Exercise D2

(a) Let  $x = 0.\overline{481}$ .

Multiplying both sides by  $10^3$ , we obtain

$$1000x = 481.\overline{481} = 481 + x.$$

Hence

$$999x = 481, \text{ so } x = 481/999 = 13/27.$$

(b) Let  $y = 0.\overline{1}$ .

Multiplying both sides by 10, we obtain

$$10y = 1.\overline{1} = 1 + y.$$

Hence

$$9y = 1, \text{ so } y = 1/9.$$

Thus

$$0.48\overline{1} = \frac{48}{100} + \frac{y}{100} = \frac{48}{100} + \frac{1}{900} = \frac{433}{900}.$$

## Solution to Additional Exercise D3

If  $x = 0.\overline{21}$ , then multiplying both sides by 100 we obtain

$$100x = 21.\overline{21} = 21 + x.$$

Hence

$$99x = 21, \text{ so } x = 21/99.$$

If  $y = 0.\overline{2}$ , then multiplying both sides by 10 we obtain

$$10y = 2.\overline{2} = 2 + y.$$

Hence

$$9y = 2, \text{ so } y = 2/9.$$

Thus

$$x + y = \frac{21}{99} + \frac{2}{9} = \frac{43}{99} = 0.\overline{43}$$

and

$$xy = \frac{21}{99} \times \frac{2}{9} = \frac{14}{297} = 0.\overline{047138}.$$

## Solution to Additional Exercise D4

For example,

$$x = 0.1195 \text{ and } y = 0.119501001\dots,$$

where  $501001\dots$  is a non-recurring tail.

## Solution to Additional Exercise D5

(a) Note that this inequality should not be solved by cross-multiplying because  $x^2 - 4$  can be positive, zero or negative depending on the value of  $x$ . Rearranging this inequality we have

$$\begin{aligned} \frac{x-1}{x^2+4} < \frac{x+1}{x^2-4} &\iff 0 < \frac{x+1}{x^2-4} - \frac{x-1}{x^2+4} \\ &\iff 0 < \frac{2x(x+4)}{(x^2-4)(x^2+4)} \\ &\iff 0 < \frac{2x(x+4)}{(x-2)(x+2)(x^2+4)}. \end{aligned}$$

Using a table of signs, including the factors

$x, x+4, x-2, x+2$  and  $x^2+4$ , we find that the solution set is

$$\left\{ x : \frac{x-1}{x^2+4} < \frac{x+1}{x^2-4} \right\} = (-\infty, -4) \cup (-2, 0) \cup (2, \infty).$$

(b) The expression  $\sqrt{4x-3}$  is defined, and non-negative, when  $4x-3 \geq 0$ ; that is, for  $x \geq \frac{3}{4}$ .

For  $x \geq \frac{3}{4}$ , we see that

$$\begin{aligned} \sqrt{4x-3} > x &\iff 4x-3 > x^2 \\ &\iff 0 > x^2 - 4x + 3 \\ &\iff 0 > (x-1)(x-3). \end{aligned}$$

Hence the solution set is

$$\{x : \sqrt{4x-3} > x\} = (1, 3).$$

(c) Rearranging this inequality we have

$$\begin{aligned} |17 - 2x^4| \leq 15 &\iff -15 \leq 17 - 2x^4 \leq 15 \\ &\iff -32 \leq -2x^4 \leq -2 \\ &\iff 16 \geq x^4 \geq 1 \\ &\iff 2 \geq |x| \geq 1. \end{aligned}$$

Hence the solution set is

$$\{x : |17 - 2x^4| \leq 15\} = [-2, -1] \cup [1, 2].$$

**Solution to Additional Exercise D6**

Suppose that  $|a| \leq 1$ . By the backwards form of the Triangle Inequality,

$$\begin{aligned} |a - 3| &\geq ||a| - 3| \\ &= 3 - |a| \quad (\text{since } |a| - 3 < 0) \\ &\geq 2 \quad (\text{since } |a| \leq 1). \end{aligned}$$

Hence

$$|a| \leq 1 \implies |a - 3| \geq 2.$$

**Solution to Additional Exercise D7**

We rearrange the inequality into a simpler equivalent form:

$$\begin{aligned} (a^2 + b^2)(c^2 + d^2) &\geq (ac + bd)^2 \\ \iff a^2c^2 + a^2d^2 + b^2c^2 + b^2d^2 \\ &\geq a^2c^2 + 2acbd + b^2d^2 \\ \iff a^2d^2 + b^2c^2 - 2acbd &\geq 0 \\ \iff (ad - bc)^2 &\geq 0. \end{aligned}$$

The last inequality is true for all  $a, b, c, d \in \mathbb{R}$ , so the first inequality must also be true for all  $a, b, c, d \in \mathbb{R}$ , as required.

**Solution to Additional Exercise D8**

(a) We substitute  $x = 2$  in the Binomial Theorem and notice that all the terms are positive; this gives, for  $n \geq 2$ ,

$$\begin{aligned} 3^n &= (1 + 2)^n \geq 1 + 2n + \frac{n(n-1)}{2!} 2^2 \\ &= 1 + 2n + 2n(n-1) \\ &= 1 + 2n^2. \end{aligned}$$

For  $n = 1$ ,

$$3^n = 3 = 1 + 2n^2.$$

Thus

$$3^n \geq 1 + 2n^2, \text{ for } n \geq 1.$$

(b) To use mathematical induction, we let  $P(n)$  be the statement

$$3^n \geq 1 + 2n^2.$$

As noted in part (a),

$$3^n = 3 = 1 + 2n^2, \text{ for } n = 1,$$

and so  $P(1)$  is true.

Now let  $k \geq 1$  and assume that  $P(k)$  is true; that is,

$$3^k \geq 2k^2 + 1.$$

We wish to deduce that  $P(k+1)$  is true; that is,

$$3^{k+1} \geq 2(k+1)^2 + 1.$$

Multiplying the inequality  $3^k \geq 2k^2 + 1$  by 3 we get

$$3^{k+1} \geq 3(2k^2 + 1),$$

so it is sufficient for our purposes to prove that

$$3(2k^2 + 1) \geq 2(k+1)^2 + 1, \text{ for } k \geq 1.$$

Now

$$\begin{aligned} 3(2k^2 + 1) &\geq 2(k+1)^2 + 1 \\ \iff 6k^2 + 3 &\geq 2k^2 + 4k + 3 \\ \iff 4k^2 &\geq 4k, \end{aligned}$$

and this last equivalent inequality is certainly true for  $k \geq 1$ . Thus

$$P(k) \implies P(k+1), \text{ for } k \geq 1.$$

Hence, by mathematical induction,  $P(n)$  is true, for  $n \geq 1$ .

**Solution to Additional Exercise D9**

Since  $x = 2/n > -1$  for  $n \geq 1$ , we can apply Bernoulli's Inequality with  $x = 2/n$  to obtain

$$\left(1 + \frac{2}{n}\right)^n \geq 1 + n \left(\frac{2}{n}\right) = 3.$$

Hence (by Rule 5 with  $p = n$ ) we have

$$1 + \frac{2}{n} \geq 3^{1/n}, \text{ for } n \geq 1.$$

Next, since  $x = -2/(3n) > -1$  for  $n \geq 1$ , we can apply Bernoulli's Inequality with  $x = -2/(3n)$  to obtain

$$\left(1 - \frac{2}{3n}\right)^n \geq 1 + n \left(-\frac{2}{3n}\right) = \frac{1}{3}.$$

Hence (by Rule 5 with  $p = n$ ) we have

$$\frac{3n-2}{3n} \geq \left(\frac{1}{3}\right)^{1/n}.$$

We can rewrite this inequality in the form

$$3^{1/n} \geq \frac{3n}{3n-2} = 1 + \frac{2}{3n-2}, \text{ for } n \geq 1.$$

Putting these two results together, we get the required inequality.

## Solution to Additional Exercise D10

Let  $P(n)$  be the statement

$$\frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-1)}{2 \cdot 4 \cdot 6 \cdot \dots \cdot (2n)} \leq \sqrt{\frac{3/4}{2n+1}}.$$

$P(1)$  is true since

$$\frac{1}{2} = \sqrt{\frac{1}{4}} = \sqrt{\frac{3/4}{2+1}}.$$

Now let  $k \geq 1$  and assume that  $P(k)$  is true; that is,

$$\frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2k-1)}{2 \cdot 4 \cdot 6 \cdot \dots \cdot (2k)} \leq \sqrt{\frac{3/4}{2k+1}}.$$

We wish to deduce that  $P(k+1)$  is true; that is,

$$\frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2(k+1)-1)}{2 \cdot 4 \cdot 6 \cdot \dots \cdot (2(k+1))} \leq \sqrt{\frac{3/4}{2(k+1)+1}}$$

which simplifies to

$$\frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2k+1)}{2 \cdot 4 \cdot 6 \cdot \dots \cdot (2k+2)} \leq \sqrt{\frac{3/4}{2k+3}}.$$

Now

$$\begin{aligned} & \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2k+1)}{2 \cdot 4 \cdot 6 \cdot \dots \cdot (2k+2)} \\ &= \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2k-1)(2k+1)}{2 \cdot 4 \cdot 6 \cdot \dots \cdot (2k)(2k+2)} \\ &\leq \sqrt{\frac{3/4}{2k+1}} \times \frac{2k+1}{2k+2}, \end{aligned}$$

using the hypothesis that  $P(k)$  is true. Hence, to prove that the statement  $P(k+1)$  holds, we need to prove that

$$\sqrt{\frac{3/4}{2k+1}} \times \frac{2k+1}{2k+2} \leq \sqrt{\frac{3/4}{2k+3}}.$$

Simplifying this inequality gives

$$\frac{\sqrt{2k+1}}{2k+2} \leq \sqrt{\frac{1}{2k+3}}.$$

But

$$\begin{aligned} \frac{\sqrt{2k+1}}{2k+2} &\leq \sqrt{\frac{1}{2k+3}} \\ \iff (2k+1)(2k+3) &\leq (2k+2)^2 \\ \iff 4k^2 + 8k + 3 &\leq 4k^2 + 8k + 4. \end{aligned}$$

This last inequality is certainly true. Thus we have shown that

$$P(k) \implies P(k+1), \text{ for } k \geq 1.$$

Hence, by mathematical induction,  $P(n)$  is true, for  $n \geq 1$ .

## Solution to Additional Exercise D11

(a) The set  $E_1$  is bounded above by 1, for example, since

$$x \leq 1, \quad \text{for } x \in E_1.$$

But  $E_1$  does not have a maximum element. For each  $x \in E_1$  we have  $0 \leq x < 1$ , so, by the Density Property, there is a rational number  $y$  such that  $x < y < 1$ . Since  $y \in E_1$ , we deduce that  $x$  is not a maximum element of  $E_1$ .

The set  $E_2$  is bounded above by 4, for example, since

$$\left(1 + \frac{1}{n}\right)^2 \leq (1+1)^2 = 4, \quad \text{for } n = 1, 2, \dots$$

Also,  $4 \in E_2$  since  $4 = \left(1 + \frac{1}{n}\right)^2$  when  $n = 1$  and so 4 is the maximum element of  $E_2$ .

(b) The set  $E_1$  is bounded below by 0, for example, since

$$x \geq 0, \quad \text{for } x \in E_1.$$

Also,  $0 \in E_1$  and so 0 is the minimum element of  $E_1$ .

The set  $E_2$  is bounded below by 1, for example, since

$$\left(1 + \frac{1}{n}\right)^2 \geq 1, \quad \text{for } n = 1, 2, \dots$$

But  $E_2$  does not have a minimum element. For each  $x$  in  $E_2$ , we have  $x = \left(1 + \frac{1}{n}\right)^2$ , for some  $n$ , so there is another element of  $E_2$ , for example  $\left(1 + \frac{1}{n+1}\right)^2$ , such that

$$\left(1 + \frac{1}{n+1}\right)^2 < \left(1 + \frac{1}{n}\right)^2 = x.$$

Hence  $x$  is not a minimum element of  $E_2$ .

(c) As noted in part (a), 1 is an upper bound of  $E_1$ . To show that  $M = 1$  is the *least* upper bound, we prove that, if  $M' < 1$ , then there is an element  $x$  in  $E_1$  which is greater than  $M'$ . Suppose that  $M' < 1$ . Then, by the Density Property, there is a rational number  $x$  such that

$$M' < x < 1 \text{ and } x \geq 0,$$

so  $x \in E_1$  and  $x > M'$ . Thus  $M'$  is not an upper bound of  $E_1$ . Hence the least upper bound of  $E_1$  is 1.

The least upper bound of  $E_2$  is 4, since  $\max E_2 = 4$ .

(d) The greatest lower bound of  $E_1$  is 0, since  $\min E_1 = 0$ .

As noted in part (b), 1 is a lower bound of  $E_2$ . To see that  $m = 1$  is the *greatest* lower bound of  $E_2$ , we have to prove that, if  $m' > 1$ , then there is an element  $(1 + 1/n)^2$  of  $E_2$  such that  $(1 + 1/n)^2 < m'$ . Suppose that  $m' > 1$ .

Since  $\sqrt{m'} > 1$ , we have

$$\begin{aligned} \left(1 + \frac{1}{n}\right)^2 < m' &\iff 1 + \frac{1}{n} < \sqrt{m'} \\ &\iff \frac{1}{n} < \sqrt{m'} - 1 \\ &\iff n > 1/(\sqrt{m'} - 1). \end{aligned}$$

By the Archimedean Property, we can choose a positive integer  $n$  so large that  $n > 1/(\sqrt{m'} - 1)$ . Hence, for this value of  $n$ , we have

$$\left(1 + \frac{1}{n}\right)^2 < m'$$

and so  $m'$  is not a lower bound of  $E_2$ . Hence the greatest lower bound of  $E_2$  is 1.

# Additional exercises for Unit D2

## Section 1

### Additional Exercise D12

Calculate the first five terms of each of the following sequences  $(a_n)$  and draw a sequence diagram in each case.

- (a)  $a_n = n^2 - 4n + 4, \quad n = 1, 2, \dots$
- (b)  $a_n = (-1)^{n+1}/n!, \quad n = 1, 2, \dots$
- (c)  $a_n = \sin(\frac{1}{4}n\pi), \quad n = 1, 2, \dots$

### Additional Exercise D13

Determine which of the following sequences  $(a_n)$  are monotonic.

- (a)  $a_n = \frac{n}{n+1}, \quad n = 1, 2, \dots$
- (b)  $a_n = \frac{(-1)^n}{n}, \quad n = 1, 2, \dots$
- (c)  $a_n = 2^{1/n}, \quad n = 1, 2, \dots$

### Additional Exercise D14

Prove that the following sequences  $(a_n)$  are each eventually monotonic.

- (a)  $a_n = 5^n/n!, \quad n = 1, 2, \dots$
- (b)  $a_n = n + 8/n, \quad n = 1, 2, \dots$

## Section 2

### Additional Exercise D15

For each of the following sequences  $(a_n)$  and positive numbers  $\varepsilon$ , find an integer  $N$  such that

$$|a_n| < \varepsilon, \quad \text{for all } n > N.$$

- (a)  $a_n = (-1)^n/n^5, \quad \varepsilon = 0.001$
- (b)  $a_n = 1/(2n+1)^2, \quad \varepsilon = 0.002$

### Additional Exercise D16

Use the definition of a null sequence to prove that the two sequences in Additional Exercise D15 are null.

### Additional Exercise D17

Prove that the following sequences  $(a_n)$  are not null.

- (a)  $a_n = \sqrt{n}, \quad n = 1, 2, \dots$
- (b)  $a_n = 1 + \frac{(-1)^n}{n}, \quad n = 1, 2, \dots$

### Additional Exercise D18

Use the fact that  $(1/n)$  is null to deduce that the following sequences  $(a_n)$  are null. State which rules you use.

- (a)  $a_n = \frac{2}{\sqrt{n}} + \frac{3}{n^7}, \quad n = 1, 2, \dots$
- (b)  $a_n = \frac{\cos n}{n^2 + 1}, \quad n = 1, 2, \dots$
- (c)  $a_n = \frac{n!}{n^n}, \quad n = 1, 2, \dots$

### Additional Exercise D19

Use the identity  $a - b = (a^2 - b^2)/(a + b)$  to prove that the sequence  $(a_n)$  is null, where

$$a_n = \sqrt{n+1} - \sqrt{n}, \quad n = 1, 2, \dots$$

### Additional Exercise D20

Use the list of basic null sequences to prove that the following sequences  $(a_n)$  are null. State which rules you use.

- (a)  $a_n = \frac{3}{4^n} + \frac{2n}{3^n}, \quad n = 1, 2, \dots$
- (b)  $a_n = \frac{6n^{10}}{n!}, \quad n = 1, 2, \dots$
- (c)  $a_n = \frac{n^{10}10^n}{n!}, \quad n = 1, 2, \dots$

*Hint:* In part (c), you can express the sequence as a product of basic null sequences.



**Additional Exercise D21 Challenging**

Determine whether or not the sequence  $(a_n)$  is null, where

$$a_n = (n+1)^{3/2} - n^{3/2}, \quad n = 1, 2, \dots$$

*Hint:* Use the same identity as in Additional Exercise D19.

**Section 3****Additional Exercise D22**

Show that the following sequences  $(a_n)$  converge to 1, by calculating  $a_n - 1$  in each case.

(a)  $a_n = \frac{n-1}{n+3}, \quad n = 1, 2, \dots$

(b)  $a_n = \frac{n^2}{n^2 + n + 1}, \quad n = 1, 2, \dots$

**Additional Exercise D23**

Use the Combination Rules to find the limits of the following sequences  $(a_n)$ .

(a)  $a_n = \frac{n^2}{n^2 + n + 1}, \quad n = 1, 2, \dots$

(b)  $a_n = \frac{n^2 - 2^n}{2^n + n^{20}}, \quad n = 1, 2, \dots$

(c)  $a_n = \frac{5n! + 5^n}{n^{100} + n!}, \quad n = 1, 2, \dots$

**Additional Exercise D24**

Prove that if  $(a_n)$  is convergent with limit  $l$ , then  $(a_n^2)$  is convergent with limit  $l^2$ .

**Section 4****Additional Exercise D25**

Classify the following sequences  $(a_n)$  as bounded or unbounded and as convergent or divergent.

(a)  $a_n = n^{1/4}, \quad n = 1, 2, \dots$

(b)  $a_n = 100^n/n!, \quad n = 1, 2, \dots$

**Additional Exercise D26**

Use the Reciprocal Rule to prove that the following sequences  $(a_n)$  tend to infinity.

(a)  $a_n = n!/n^3, \quad n = 1, 2, \dots$

(b)  $a_n = n^2 + 2n, \quad n = 1, 2, \dots$

(c)  $a_n = n^2 - 2n, \quad n = 1, 2, \dots$

(d)  $a_n = n! - n^3 - 3^n, \quad n = 1, 2, \dots$

**Additional Exercise D27**

Use the Subsequence Rules to prove that the following sequences  $(a_n)$  are divergent.


(a)  $a_n = (-1)^n 2^n, \quad n = 1, 2, \dots$

(b)  $a_n = \frac{(-1)^n n^2}{2n^2 + 1}, \quad n = 1, 2, \dots$


# Solutions to additional exercises for Unit D2

## Solution to Additional Exercise D12


(a) 1, 0, 1, 4, 9.



(b)  $1, -\frac{1}{2}, \frac{1}{6}, -\frac{1}{24}, \frac{1}{120}$ .



(c)  $\frac{1}{\sqrt{2}}, 1, \frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}}$ .



## Solution to Additional Exercise D13

(a) Since  $a_n > 0$  for all  $n$ , we can use Strategy D4. We have

$$a_n = \frac{n}{n+1} \text{ and } a_{n+1} = \frac{n+1}{n+2}$$

so

$$\begin{aligned} \frac{a_{n+1}}{a_n} &= \frac{(n+1)/(n+2)}{n/(n+1)} \\ &= \frac{n^2 + 2n + 1}{n^2 + 2n} \geq 1, \quad \text{for } n = 1, 2, \dots \end{aligned}$$

Thus  $(a_n)$  is increasing, so  $(a_n)$  is monotonic.

(b) We have

$$a_1 = -1, \quad a_2 = \frac{1}{2}, \quad a_3 = -\frac{1}{3}.$$

Since  $a_1 < a_2$ , we know that  $(a_n)$  is not decreasing; and since  $a_3 < a_2$ , we know that  $(a_n)$  is not increasing.

Hence  $(a_n)$  is not monotonic.

(c) We have

$$a_{n+1} = 2^{1/(n+1)} \leq 2^{1/n} = a_n, \quad \text{for } n = 1, 2, \dots$$

This holds because, by the rules for inequalities (Rule 5 with  $p = n(n+1)$ ),

$$2^{1/(n+1)} \leq 2^{1/n} \iff 2^n \leq 2^{n+1},$$

and the right-hand inequality is true for  $n = 1, 2, \dots$

Thus  $(a_n)$  is decreasing, so  $(a_n)$  is monotonic.

## Solution to Additional Exercise D14

(a) We have  $a_n > 0$  for all  $n$  and

$$\frac{a_{n+1}}{a_n} = \frac{5^{n+1}/(n+1)!}{5^n/n!} = \frac{5}{n+1}.$$

Now

$$\frac{5}{n+1} \leq 1, \quad \text{for } n = 4, 5, 6, \dots,$$

so

$$a_{n+1} \leq a_n, \quad \text{for } n = 4, 5, 6, \dots$$

Hence  $(a_n)$  is eventually decreasing, so  $(a_n)$  is eventually monotonic.

(b) We have

$$\begin{aligned} a_{n+1} - a_n &= \left(n+1 + \frac{8}{n+1}\right) - \left(n + \frac{8}{n}\right) \\ &= 1 - \frac{8}{n(n+1)}. \end{aligned}$$

Now

$$\begin{aligned} 1 - \frac{8}{n(n+1)} \geq 0 &\iff \frac{8}{n(n+1)} \leq 1 \\ &\iff n(n+1) \geq 8 \end{aligned}$$

and this final inequality holds for  $n \geq 3$ . So

$$a_{n+1} \geq a_n, \quad \text{for } n \geq 3.$$

Hence  $(a_n)$  is eventually increasing, so  $(a_n)$  is eventually monotonic.

## Solution to Additional Exercise D15

(a) We have  $|a_n| = \left|\frac{(-1)^n}{n^5}\right| = \frac{1}{n^5}$ , and

$$\begin{aligned} \frac{1}{n^5} < 0.001 &\iff n^5 > 1000 \\ &\iff n > 1000^{1/5} \approx 3.98. \end{aligned}$$

Hence

$$|a_n| < 0.001, \quad \text{for all } n > 3,$$

so we can take  $N = 3$ .

(b) We have  $|a_n| = 1/(2n+1)^2$ , and

$$\begin{aligned} \frac{1}{(2n+1)^2} < 0.002 &\iff (2n+1)^2 > 500 \\ &\iff n > \frac{1}{2}(\sqrt{500} - 1) \approx 10.68. \end{aligned}$$

Hence

$$|a_n| < 0.002, \quad \text{for all } n > 10,$$

so we can take  $N = 10$ .

### Solution to Additional Exercise D16

(a) We want to show that:

for each  $\varepsilon > 0$ , there is an integer  $N$  such that

$$\left| \frac{(-1)^n}{n^5} \right| = \frac{1}{n^5} < \varepsilon, \quad \text{for all } n > N. \quad (\text{S1})$$

We know that

$$\begin{aligned} \frac{1}{n^5} < \varepsilon &\iff n^5 > \frac{1}{\varepsilon} \\ &\iff n > \left( \frac{1}{\varepsilon} \right)^{1/5}. \end{aligned}$$

If we take  $N \geq (1/\varepsilon)^{1/5}$ , then statement (S1) holds. Hence  $(a_n)$  is null.

(b) We want to show that:

for each  $\varepsilon > 0$ , there is an integer  $N$  such that

$$\frac{1}{(2n+1)^2} < \varepsilon, \quad \text{for all } n > N. \quad (\text{S2})$$

We know that

$$\begin{aligned} \frac{1}{(2n+1)^2} < \varepsilon &\iff (2n+1)^2 > 1/\varepsilon \\ &\iff n > \frac{1}{2} \left( \sqrt{1/\varepsilon} - 1 \right). \end{aligned}$$

If we take  $N \geq \frac{1}{2}(\sqrt{1/\varepsilon} - 1)$ , then statement (S2) holds. Hence  $(a_n)$  is null.

### Solution to Additional Exercise D17

(a) We use Strategy D5. We have to find a positive number  $\varepsilon$  for which there is no integer  $N$  such that

$$|a_n| = \sqrt{n} < \varepsilon, \quad \text{for all } n > N.$$

Since  $\sqrt{n} > \frac{1}{2}$  for all  $n$ , we can take  $\varepsilon = \frac{1}{2}$ , for example. Hence  $(a_n)$  is not null.

(b) We use Strategy D5. We have to find a positive number  $\varepsilon$  for which there is no integer  $N$  such that

$$|a_n| = \left| 1 + \frac{(-1)^n}{n} \right| < \varepsilon, \quad \text{for all } n > N.$$

Since

$$\left| \frac{(-1)^n}{n} \right| = \frac{1}{n} \leq \frac{1}{2} \quad \text{for } n \geq 2,$$

we have

$$-\frac{1}{2} \leq \frac{(-1)^n}{n} \leq \frac{1}{2}, \quad \text{for } n \geq 2,$$

so

$$\frac{1}{2} \leq 1 + \frac{(-1)^n}{n} \leq \frac{3}{2}, \quad \text{for } n \geq 2.$$

So we can take  $\varepsilon = \frac{1}{2}$ , for example. Hence  $(a_n)$  is not null.

### Solution to Additional Exercise D18

(a) The sequences  $\left( \frac{2}{\sqrt{n}} \right)$  and  $\left( \frac{3}{n^7} \right)$  are null, by the Power Rule and the Multiple Rule.

Hence  $(a_n)$  is null, by the Sum Rule.

(b) We guess that  $(a_n)$  is dominated by  $(b_n)$ , where

$$b_n = \frac{1}{n}, \quad n = 1, 2, \dots$$

This holds because

$$|\cos n| \leq 1 \quad \text{for } n = 1, 2, \dots,$$

and

$$n^2 + 1 \geq n \quad \text{for } n = 1, 2, \dots,$$

so that

$$\left| \frac{\cos n}{n^2 + 1} \right| \leq \frac{1}{n}, \quad \text{for } n = 1, 2, \dots$$

Since  $(b_n)$  is null, we deduce that  $(a_n)$  is null, by the Squeeze Rule.

(c) Since

$$\frac{n!}{n^n} = \frac{1 \times 2 \times \dots \times n}{n \times n \times \dots \times n} = \frac{1}{n} \times \frac{2}{n} \times \dots \times \frac{n-1}{n} \times \frac{n}{n},$$

we have

$$0 \leq \frac{n!}{n^n} \leq \frac{1}{n}, \quad \text{for } n = 1, 2, \dots$$

Hence  $(a_n)$  is null, by the Squeeze Rule.

### Solution to Additional Exercise D19

Using the given identity, we obtain

$$\sqrt{n+1} - \sqrt{n} = \frac{(n+1) - n}{\sqrt{n+1} + \sqrt{n}} = \frac{1}{\sqrt{n+1} + \sqrt{n}}.$$

Thus, for  $n = 1, 2, \dots$ , we have

$$0 \leq \sqrt{n+1} - \sqrt{n} = \frac{1}{\sqrt{n+1} + \sqrt{n}} < \frac{1}{\sqrt{n}}.$$

Now the sequence  $\left(\frac{1}{\sqrt{n}}\right)$  is null, by the

Power Rule.

Hence  $(a_n)$  is null, by the Squeeze Rule.

### Solution to Additional Exercise D20

(a) Since

$$\frac{1}{4^n} = \left(\frac{1}{4}\right)^n \quad \text{and} \quad \frac{n}{3^n} = n \left(\frac{1}{3}\right)^n,$$

both  $(1/4^n)$  and  $(n/3^n)$  are basic null sequences, so  $(a_n)$  is null, by the Multiple Rule and the Sum Rule.

(b)  $(n^{10}/n!)$  is a basic null sequence, so  $(a_n)$  is a null sequence, by the Multiple Rule.

(c) First note that

$$\frac{n^{10}10^n}{n!} = \frac{n^{10}}{10^n} \frac{100^n}{n!}, \quad \text{for } n = 1, 2, \dots$$

Since  $(n^{10}/10^n)$  and  $(100^n/n!)$  are basic null sequences,  $(a_n)$  is null, by the Product Rule.

### Solution to Additional Exercise D21

The sequence  $(a_n)$  is not null.

To prove this, we use Strategy D5. We have to find a positive number  $\varepsilon$  for which there is no integer  $N$  such that

$$|a_n| < \varepsilon, \quad \text{for all } n > N.$$

Using the same identity as in

Additional Exercise D19, we have that

$$\begin{aligned} a_n &= (n+1)^{3/2} - n^{3/2} = \frac{(n+1)^3 - n^3}{(n+1)^{3/2} + n^{3/2}} \\ &= \frac{3n^2 + 3n + 1}{(n+1)^{3/2} + n^{3/2}} \\ &> \frac{3n^2}{(n+1)^{3/2} + n^{3/2}} \\ &\geq \frac{3n^2}{(2n)^{3/2} + n^{3/2}} \\ &= \frac{3n^2}{(2^{3/2} + 1)n^{3/2}} > \frac{3}{4}\sqrt{n}. \end{aligned}$$

In particular,  $a_n > \frac{3}{4}\sqrt{n} > \frac{3}{4}$ , for  $n = 1, 2, \dots$

So we can take  $\varepsilon = \frac{3}{4}$ , for example.

### Solution to Additional Exercise D22

(a) We have

$$a_n - 1 = \frac{n-1}{n+3} - 1 = \frac{-4}{n+3}.$$

Now  $\left(\frac{-4}{n+3}\right)$  is a null sequence, by the Squeeze Rule, since

$$\left| \frac{-4}{n+3} \right| = \frac{4}{n+3} \leq \frac{4}{n}, \quad \text{for } n = 1, 2, \dots$$

Thus  $(a_n)$  is convergent with limit 1.

(b) We have

$$a_n - 1 = \frac{n^2}{n^2 + n + 1} - 1 = -\frac{n+1}{n^2 + n + 1}.$$

Now  $\left(-\frac{n+1}{n^2 + n + 1}\right)$  is a null sequence, by the Squeeze Rule, since

$$\left| -\frac{n+1}{n^2 + n + 1} \right| \frac{n+1}{n^2 + n + 1} = \frac{n+1}{n(n+1)+1} \leq \frac{1}{n},$$

for  $n = 1, 2, \dots$ . Thus  $(a_n)$  is convergent with limit 1.

### Solution to Additional Exercise D23

In each case we apply Strategy D7.

(a) The dominant term is  $n^2$ , so we write

$$a_n = \frac{n^2}{n^2 + n + 1} = \frac{1}{1 + 1/n + 1/n^2}.$$

Since  $(1/n)$  and  $(1/n^2)$  are basic null sequences, we find, by the Combination Rules, that

$$\lim_{n \rightarrow \infty} a_n = \frac{1}{1 + 0 + 0} = 1.$$

(b) The dominant term is  $2^n$ , so we write

$$\frac{n^2 - 2^n}{2^n + n^{20}} = \frac{n^2/2^n - 1}{1 + n^{20}/2^n}.$$

Since  $(n^2/2^n) = (n^2(\frac{1}{2})^n)$  and  $(n^{20}/2^n) = (n^{20}(\frac{1}{2})^n)$  are basic null sequences, we find, by the Combination Rules, that

$$\lim_{n \rightarrow \infty} a_n = \frac{0 - 1}{1 + 0} = -1.$$

(c) The dominant term is  $n!$ , so we write

$$\frac{5n! + 5^n}{n^{100} + n!} = \frac{5 + 5^n/n!}{n^{100}/n! + 1}.$$

Since  $(5^n/n!)$  and  $(n^{100}/n!)$  are basic null sequences, we find, by the Combination Rules, that

$$\lim_{n \rightarrow \infty} a_n = \frac{5 + 0}{0 + 1} = 5.$$

### Solution to Additional Exercise D24

The result follows from the Product Rule with  $b_n = a_n$ , for  $n = 1, 2, \dots$ .

### Solution to Additional Exercise D25

(a) This sequence is unbounded and hence divergent, by Corollary D15.

(b) This sequence is a basic null sequence. Hence it is convergent and therefore bounded, by Theorem D14.

### Solution to Additional Exercise D26

(a) Each term of the sequence  $(a_n)$  is positive. Also

$$\frac{1}{a_n} = \frac{n^3}{n!}$$

and  $(n^3/n!)$  is a basic null sequence.

Hence  $a_n \rightarrow \infty$ , by the Reciprocal Rule.

(b) Each term of the sequence  $(a_n)$  is positive and

$$0 \leq \frac{1}{a_n} = \frac{1}{n^2 + 2n} \leq \frac{1}{n^2}, \quad \text{for } n = 1, 2, \dots$$

Since  $(1/n^2)$  is null, we deduce that  $(1/a_n)$  is null, by the Squeeze Rule.

Hence  $a_n \rightarrow \infty$ , by the Reciprocal Rule.

(c) The dominant term is  $n^2$ , so we first write

$$a_n = n^2 - 2n = n^2(1 - 2/n), \quad \text{for } n = 1, 2, \dots$$

Since  $2/n < 1$ , for  $n > 2$ , we deduce that  $(a_n)$  is eventually positive.

Next we write

$$\frac{1}{a_n} = \frac{1}{n^2 - 2n} = \frac{1/n^2}{1 - 2/n}.$$

Now  $(1/n)$  and  $(1/n^2)$  are basic null sequences. Thus, by the Combination Rules,

$$\lim_{n \rightarrow \infty} \frac{1}{a_n} = \frac{0}{1 - 0} = 0.$$

Hence  $a_n \rightarrow \infty$ , by the Reciprocal Rule.

(d) The dominant term is  $n!$ , so we first write

$$a_n = n! - n^3 - 3^n = n! \left( 1 - \frac{n^3}{n!} - \frac{3^n}{n!} \right),$$

for  $n = 1, 2, \dots$

Since  $(n^3/n!)$  and  $(3^n/n!)$  are basic null sequences, we know that there is some integer  $N$  such that

$$\frac{n^3}{n!} + \frac{3^n}{n!} < 1, \quad \text{for } n > N.$$

Thus  $(a_n)$  is eventually positive.

Next we write

$$\frac{1}{a_n} = \frac{1}{n! - n^3 - 3^n} = \frac{1/n!}{1 - n^3/n! - 3^n/n!}.$$

Now  $(1/n!)$ ,  $(n^3/n!)$  and  $(3^n/n!)$  are basic null sequences. Thus, by the Combination Rules,

$$\lim_{n \rightarrow \infty} \frac{1}{a_n} = \frac{0}{1 - 0 - 0} = 0.$$

Hence  $a_n \rightarrow \infty$ , by the Reciprocal Rule.

### Solution to Additional Exercise D27

(a) We have

$$a_{2k} = 2^{2k} = 4^k, \quad \text{for } k = 1, 2, \dots$$

Each term of  $(a_{2k})$  is positive. Also

$$\frac{1}{a_{2k}} = \frac{1}{4^k} = \left( \frac{1}{4} \right)^k.$$

and  $((1/4)^k)$  is a basic null sequence. Hence  $a_{2k} \rightarrow \infty$ , by the Reciprocal Rule. Thus  $(a_n)$  is divergent, by the Second Subsequence Rule.

(b) We have

$$a_{2k} = \frac{(2k)^2}{2(2k)^2 + 1} = \frac{4k^2}{8k^2 + 1} = \frac{4}{8 + 1/k^2}$$

and

$$\begin{aligned} a_{2k-1} &= \frac{-(2k-1)^2}{2(2k-1)^2 + 1} = -\frac{4k^2 - 4k + 1}{8k^2 - 8k + 3} \\ &= -\frac{4 - 4/k + 1/k^2}{8 - 8/k + 3/k^2}. \end{aligned}$$

Now  $(1/k)$  and  $(1/k^2)$  are basic null sequences, so

$$\lim_{k \rightarrow \infty} a_{2k} = \frac{1}{2}, \quad \text{whereas} \quad \lim_{k \rightarrow \infty} a_{2k-1} = -\frac{1}{2},$$

by the Combination Rules.

Hence  $(a_n)$  is divergent, by the First Subsequence Rule.

# Additional exercises for Unit D3

## Section 1

### Additional Exercise D28

Prove that  $\sum_{n=0}^{\infty} \left(-\frac{4}{5}\right)^n$  is convergent and find its sum.

### Additional Exercise D29

Prove that

$$\frac{1}{2n-1} - \frac{1}{2n+1} = \frac{2}{4n^2-1}, \quad \text{for } n = 1, 2, \dots,$$

and deduce that

$$\sum_{n=1}^{\infty} \frac{1}{4n^2-1} = \frac{1}{2}.$$

### Additional Exercise D30

Determine the behaviour of the following series. If the series is convergent, find its sum.

(a)  $\sum_{n=1}^{\infty} \left( \left(\frac{4}{5}\right)^n + \frac{4}{n(n+2)} \right)$

(b)  $\sum_{n=1}^{\infty} \left( 1 + \left(\frac{1}{2}\right)^n \right)$

*Hint:* For part (a), you may find it helpful to refer to the solution to Exercise D43 in Unit D3.

## Section 2

### Additional Exercise D31

Determine whether or not the following series are convergent.

(a)  $\sum_{n=1}^{\infty} \frac{\cos(1/n)}{2n^2+3}$  (b)  $\sum_{n=1}^{\infty} \frac{n^2}{2n^3-n}$

(c)  $\sum_{n=1}^{\infty} \frac{\sqrt{2n}}{4n^3+n+2}$  (d)  $\sum_{n=1}^{\infty} \frac{(n+1)^5}{2^n}$

(e)  $\sum_{n=1}^{\infty} \frac{n^2 3^n}{n!}$  (f)  $\sum_{n=1}^{\infty} \frac{(n!)^2}{(2n)!}$

### Additional Exercise D32 Challenging

(a) Use the Ratio Test to show that

$$\sum_{n=1}^{\infty} \frac{2^n n!}{n^n} \text{ converges}$$

but

$$\sum_{n=1}^{\infty} \frac{3^n n!}{n^n} \text{ diverges.}$$

(b) For which positive values of  $c$  can you use the Ratio Test to prove that  $\sum_{n=1}^{\infty} \frac{c^n n!}{n^n}$  is convergent?

*Hint:* Use the fact (from Unit D2) that  $(1 + 1/n)^n \rightarrow e$  as  $n \rightarrow \infty$ .

## Section 3

### Additional Exercise D33

Determine which of the following series are convergent.

(a)  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{1+\sqrt{n}}$  (b)  $\sum_{n=1}^{\infty} \frac{\sin n}{n^2}$

(c)  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} n!}{n^4+3}$  (d)  $\sum_{n=1}^{\infty} \frac{n+2^n}{3^n+5}$

(e)  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} n}{n^2+2}$

### Additional Exercise D34 Challenging

Prove that

$$\frac{1}{3n-2} + \frac{1}{3n-1} - \frac{1}{3n} > \frac{1}{3n}, \quad \text{for } n = 1, 2, \dots,$$

and deduce that the series

$$1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \dots \quad (1)$$

is divergent.

## Solutions to additional exercises for Unit D3

### Solution to Additional Exercise D28

This is a geometric series with  $a = 1$  and  $r = -\frac{4}{5}$ . Since  $|r| = |-\frac{4}{5}| = \frac{4}{5} < 1$ , the series is convergent, with sum

$$\frac{1}{1 - (-\frac{4}{5})} = \frac{5}{9}.$$

### Solution to Additional Exercise D29

For  $n = 1, 2, \dots$ ,

$$\begin{aligned} \frac{1}{2n-1} - \frac{1}{2n+1} &= \frac{(2n+1) - (2n-1)}{(2n-1)(2n+1)} \\ &= \frac{2}{(2n-1)(2n+1)} = \frac{2}{4n^2-1}, \end{aligned}$$

as required.

Thus the given series is a telescoping series and the  $n$ th partial sum  $s_n$  is given by

$$\begin{aligned} s_n &= \sum_{k=1}^n \frac{1}{4k^2-1} = \frac{1}{2} \sum_{k=1}^n \left( \frac{1}{2k-1} - \frac{1}{2k+1} \right) \\ &= \frac{1}{2} \left( \left( \frac{1}{1} - \frac{1}{3} \right) + \left( \frac{1}{3} - \frac{1}{5} \right) + \dots \right. \\ &\quad \left. + \left( \frac{1}{2n-1} - \frac{1}{2n+1} \right) \right). \end{aligned}$$

All the terms cancel out except for the first term in the first bracket and the second term in the last bracket. Thus

$$s_n = \frac{1}{2} \left( 1 - \frac{1}{2n+1} \right).$$

Since  $(1/(2n+1))$  is a null sequence, it follows that

$$\lim_{n \rightarrow \infty} s_n = \frac{1}{2},$$

so the series is convergent, with sum  $\frac{1}{2}$ .

### Solution to Additional Exercise D30

(a) The series  $\sum_{n=1}^{\infty} (\frac{4}{5})^n$  is a geometric series, with  $a = r = \frac{4}{5}$ . Since  $|r| = \frac{4}{5} < 1$ , the series is convergent, with sum

$$\frac{\frac{4}{5}}{1 - \frac{4}{5}} = 4.$$

The series  $\sum_{n=1}^{\infty} \frac{1}{n(n+2)}$  is convergent, with sum  $\frac{3}{4}$ ; see the solution to Exercise D43 in Unit D3.

Hence, by the Combination Rules,

$$\sum_{n=1}^{\infty} \left( \left( \frac{4}{5} \right)^n + \frac{4}{n(n+2)} \right) \text{ is convergent,}$$

with sum  $4 + (4 \times \frac{3}{4}) = 7$ .

(b) Let  $a_n = 1 + (\frac{1}{2})^n$ . Since  $((\frac{1}{2})^n)$  is a basic null sequence, it follows from the Combination Rules for sequences that

$$\lim_{n \rightarrow \infty} a_n = 1 \neq 0.$$

Hence, by the Non-null Test,

$$\sum_{n=1}^{\infty} (1 + (\frac{1}{2})^n) \text{ is divergent.}$$

### Solution to Additional Exercise D31

(a) We guess that this series is dominated by  $\sum (1/n^2)$ . Indeed, since

$$0 \leq 1/n \leq \frac{\pi}{2}, \quad \text{for } n = 1, 2, \dots,$$

we have

$$0 \leq \cos(1/n) \leq 1, \quad \text{for } n = 1, 2, \dots$$

Hence, for  $n = 1, 2, \dots$ ,

$$0 \leq \frac{\cos(1/n)}{2n^2+3} \leq \frac{1}{2n^2+3} \leq \frac{1}{n^2}.$$

Since  $\sum_{n=1}^{\infty} 1/n^2$  is a basic convergent series, we deduce by the Comparison Test that

$$\sum_{n=1}^{\infty} \frac{\cos(1/n)}{2n^2+3} \text{ is convergent.}$$

(b) Let

$$a_n = \frac{n^2}{2n^3 - n}, \quad \text{for } n = 1, 2, \dots$$

We guess that, for large  $n$ , the terms of this series behave like  $n^2/n^3 = 1/n$  and we know that  $\sum_{n=1}^{\infty} 1/n$  is divergent. We cannot compare  $a_n$  and  $1/n$  directly so we use the Limit Comparison Test with

$$b_n = \frac{1}{n}, \quad \text{for } n = 1, 2, \dots$$



Both  $a_n$  and  $b_n$  are positive and, by the Combination Rules for sequences,

$$\begin{aligned}\frac{a_n}{b_n} &= \left( \frac{n^2}{2n^3 - n} \right) \left( \frac{n}{1} \right) \\ &= \frac{1}{2 - 1/n^2} \rightarrow \frac{1}{2} \neq 0.\end{aligned}$$

Since  $\sum_{n=1}^{\infty} 1/n$  is a basic divergent series, we deduce by the Limit Comparison Test that

$$\sum_{n=1}^{\infty} \frac{n^2}{2n^3 - n} \text{ is divergent.}$$

(c) Let

$$a_n = \frac{\sqrt{2n}}{4n^3 + n + 2}, \quad \text{for } n = 1, 2, \dots$$

We guess that the terms of this series behave like  $\sqrt{n}/n^3 = 1/n^{5/2}$  and we know that  $\sum_{n=1}^{\infty} 1/n^{5/2}$  is convergent. We cannot compare  $a_n$  and  $1/n^{5/2}$  directly so we use the Limit Comparison Test with

$$b_n = \frac{1}{n^{5/2}}, \quad \text{for } n = 1, 2, \dots$$

Both  $a_n$  and  $b_n$  are positive and, by the Combination Rules for sequences,

$$\begin{aligned}\frac{a_n}{b_n} &= \left( \frac{\sqrt{2n}}{4n^3 + n + 2} \right) \left( \frac{n^{5/2}}{1} \right) \\ &= \frac{\sqrt{2}n^3}{4n^3 + n + 2} \\ &= \frac{\sqrt{2}}{4 + 1/n^2 + 2/n^3} \rightarrow \frac{1}{4}\sqrt{2} \neq 0.\end{aligned}$$

Since  $\sum_{n=1}^{\infty} 1/n^{5/2}$  is a basic convergent series, we deduce by the Limit Comparison Test that

$$\sum_{n=1}^{\infty} \frac{\sqrt{2n}}{4n^3 + n + 2} \text{ is convergent.}$$

(d) Let

$$a_n = \frac{(n+1)^5}{2^n}, \quad \text{for } n = 1, 2, \dots$$

We guess that the terms of this series behave like  $n^5/2^n$  and we know that  $\sum_{n=1}^{\infty} n^5/2^n$  is convergent, so we use the Limit Comparison Test with

$$b_n = \frac{n^5}{2^n}, \quad n = 1, 2, \dots$$

Both  $a_n$  and  $b_n$  are positive and, by the Combination Rules for sequences,

$$\begin{aligned}\frac{a_n}{b_n} &= \left( \frac{(n+1)^5}{2^n} \right) \left( \frac{2^n}{n^5} \right) \\ &= \left( \frac{n+1}{n} \right)^5 = \left( 1 + \frac{1}{n} \right)^5 \rightarrow 1 \neq 0.\end{aligned}$$

Since  $\sum_{n=1}^{\infty} n^5/2^n$  is a basic convergent series, we deduce by the Limit Comparison Test that

$$\sum_{n=1}^{\infty} \frac{(n+1)^5}{2^n} \text{ is convergent.}$$

(Alternatively, we can either note that

$$\sum_{n=1}^{\infty} \frac{(n+1)^5}{2^n} = \sum_{n=2}^{\infty} \frac{n^5}{2^{n-1}} = 2 \sum_{n=2}^{\infty} \frac{n^5}{2^n},$$

and use the Multiple Rule, or we can use the Ratio Test:

$$\frac{a_{n+1}}{a_n} = \left( \frac{(n+2)^5}{2^{n+1}} \right) \left( \frac{2^n}{(n+1)^5} \right) = \frac{1}{2} \left( \frac{1+2/n}{1+1/n} \right)^5,$$

which converges to  $\frac{1}{2} < 1$ .)

(e) There is no obvious basic series to compare with, so we try the Ratio Test. Let

$$a_n = \frac{n^2 3^n}{n!}, \quad \text{for } n = 1, 2, \dots$$

Then  $a_n$  is positive and

$$\begin{aligned}\frac{a_{n+1}}{a_n} &= \left( \frac{(n+1)^2 3^{n+1}}{(n+1)!} \right) \left( \frac{n!}{n^2 3^n} \right) \\ &= \frac{3(n+1)}{n^2} \\ &= 3 \left( \frac{1}{n} + \frac{1}{n^2} \right).\end{aligned}$$

Hence, by the Combination Rules for sequences,

$$\frac{a_{n+1}}{a_n} \rightarrow 0 < 1, \quad \text{as } n \rightarrow \infty.$$

Thus, by the Ratio Test,

$$\sum_{n=1}^{\infty} \frac{n^2 3^n}{n!} \text{ is convergent.}$$

(f) Again, we try the Ratio Test. Let

$$a_n = \frac{(n!)^2}{(2n)!}, \quad n = 1, 2, \dots$$

Then

$$\begin{aligned}\frac{a_{n+1}}{a_n} &= \left( \frac{((n+1)!)^2}{(2n+2)!} \right) \left( \frac{(2n)!}{(n!)^2} \right) \\ &= \frac{(n+1)^2}{(2n+2)(2n+1)} \\ &= \frac{n+1}{2(2n+1)} = \frac{1+1/n}{2(2+1/n)}.\end{aligned}$$

Hence, by the Combination Rules for sequences,

$$\frac{a_{n+1}}{a_n} \rightarrow \frac{1}{4} < 1 \quad \text{as } n \rightarrow \infty.$$

Thus, by the Ratio Test,

$$\sum_{n=1}^{\infty} \frac{(n!)^2}{(2n)!} \text{ is convergent.}$$

### Solution to Additional Exercise D32

(a) Let

$$a_n = \frac{2^n n!}{n^n}, \quad \text{for } n = 1, 2, \dots;$$

then

$$\begin{aligned} \frac{a_{n+1}}{a_n} &= \left( \frac{2^{n+1}(n+1)!}{(n+1)^{n+1}} \right) \left( \frac{n^n}{2^n n!} \right) \\ &= \frac{2(n+1)n^n}{(n+1)^{n+1}} \\ &= \frac{2n^n}{(n+1)^n} = \frac{2}{(1+1/n)^n}. \end{aligned}$$

Thus

$$\frac{a_{n+1}}{a_n} \rightarrow \frac{2}{e} < 1, \quad \text{as } n \rightarrow \infty.$$

Thus, by the Ratio Test,

$$\sum_{n=1}^{\infty} \frac{2^n n!}{n^n} \text{ is convergent.}$$

If  $a_n = 3^n n! / n^n$ , then a similar calculation yields

$$\frac{a_{n+1}}{a_n} \rightarrow \frac{3}{e} > 1, \quad \text{as } n \rightarrow \infty.$$

so, by the Ratio Test,

$$\sum_{n=1}^{\infty} \frac{3^n n!}{n^n} \text{ is divergent.}$$

(b) We follow the same approach as in part (a).

So, for any positive number  $c$ , let

$$a_n = \frac{c^n n!}{n^n}, \quad \text{for } n = 1, 2, \dots;$$

then

$$\begin{aligned} \frac{a_{n+1}}{a_n} &= \left( \frac{c^{n+1}(n+1)!}{(n+1)^{n+1}} \right) \times \left( \frac{n^n}{c^n n!} \right) \\ &= \frac{c(n+1)n^n}{(n+1)^{n+1}} \\ &= \frac{cn^n}{(n+1)^n} = \frac{c}{(1+1/n)^n}. \end{aligned}$$

Hence

$$\frac{a_{n+1}}{a_n} \rightarrow \frac{c}{e}, \quad \text{as } n \rightarrow \infty.$$

It follows, by the Ratio Test, that the series

$$\sum_{n=1}^{\infty} \frac{c^n n!}{n^n} \text{ is convergent if } 0 < c < e \text{ and divergent if}$$

$c > e$ . However, if  $c = e$  the ratio

$$\frac{a_{n+1}}{a_n} \rightarrow 1, \quad \text{as } n \rightarrow \infty;$$

and so the Ratio Test gives us no conclusion either way.

*Remark* If  $c = e$ , then the series is also divergent, but this is harder to prove. One method is to use the inequality

$$n! > \left( \frac{n+1}{e} \right)^n, \quad \text{for } n = 1, 2, \dots$$

From this inequality, we deduce that

$$\frac{e^n n!}{n^n} > \left( \frac{n+1}{n} \right)^n = (1 + 1/n)^n, \quad \text{for } n = 1, 2, \dots$$

Since  $\lim_{n \rightarrow \infty} (1 + 1/n)^n = e \neq 0$ , we deduce that  $(e^n n! / n^n)$  is not a null sequence.

Hence, by the Non-null Test,

$$\sum_{n=1}^{\infty} \frac{e^n n!}{n^n} \text{ is divergent.}$$

### Solution to Additional Exercise D33

(a) We use the Alternating Test.

Let

$$a_n = \frac{(-1)^{n+1}}{1 + \sqrt{n}}, \quad \text{for } n = 1, 2, \dots$$

Then  $a_n = (-1)^{n+1} b_n$  where

$$b_n = \frac{1}{1 + \sqrt{n}}, \quad \text{for } n = 1, 2, \dots$$

Now

1.  $b_n \geq 0$ , for  $n = 1, 2, \dots$
2.  $(b_n)$  is a null sequence by the Squeeze Rule, since

$$0 \leq b_n = \frac{1}{1 + \sqrt{n}} < \frac{1}{\sqrt{n}}, \quad \text{for } n = 1, 2, \dots$$

3.  $(b_n)$  is decreasing, because  $(1/b_n) = (1 + \sqrt{n})$  is increasing.

Hence, by the Alternating Test,

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{1 + \sqrt{n}} \text{ is convergent.}$$

(b) We guess that this series is dominated by the basic convergent series  $\sum (1/n^2)$ . Indeed, let

$$a_n = \frac{\sin n}{n^2}, \quad \text{for } n = 1, 2, \dots$$

Then

$$|a_n| = \frac{|\sin n|}{n^2} \leq \frac{1}{n^2}, \text{ for } n = 1, 2, \dots$$

Since  $\sum_{n=1}^{\infty} (1/n^2)$  is convergent, it follows from the Comparison Test that  $\sum_{n=1}^{\infty} |a_n|$  is convergent. Hence, by the Absolute Convergence Test,

$$\sum_{n=1}^{\infty} \frac{\sin n}{n^2} \text{ is convergent.}$$

(c) Let

$$a_n = \frac{(-1)^{n+1}n!}{n^4 + 3}, \text{ for } n = 1, 2, \dots$$

Then

$$\frac{1}{|a_n|} = \frac{n^4 + 3}{n!} = \frac{n^4}{n!} + \frac{3}{n!} \rightarrow 0.$$

It follows from the Reciprocal Rule that  $|a_n| \rightarrow \infty$  as  $n \rightarrow \infty$ . Thus, by the Non-null Test,

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}n!}{n^4 + 3} \text{ is divergent.}$$

(d) Let

$$a_n = \frac{n + 2^n}{3^n + 5}, \text{ for } n = 1, 2, \dots$$

We guess that  $a_n$  behaves like  $2^n/3^n = (\frac{2}{3})^n$  for large  $n$ , so we use the Limit Comparison Test with

$$b_n = \left(\frac{2}{3}\right)^n, \text{ for } n = 1, 2, \dots$$

Both  $a_n$  and  $b_n$  are positive and

$$\begin{aligned} \frac{a_n}{b_n} &= \frac{\left(\frac{n + 2^n}{3^n + 5}\right)}{\left(\frac{2}{3}\right)^n} \\ &= \frac{3^n(n + 2^n)}{2^n(3^n + 5)} \\ &= \frac{n/2^n + 1}{1 + 5/3^n} \rightarrow 1 \neq 0. \end{aligned}$$

Since  $\sum_{n=1}^{\infty} \left(\frac{2}{3}\right)^n$  is a basic convergent series, we deduce by the Limit Comparison Test that

$$\sum_{n=1}^{\infty} \frac{n + 2^n}{3^n + 5} \text{ is convergent.}$$

(e) We use the Alternating Test.

Let

$$a_n = \frac{(-1)^{n+1}n}{n^2 + 2}, \text{ for } n = 1, 2, \dots$$

Then  $a_n = (-1)^{n+1}b_n$ , where

$$b_n = \frac{n}{n^2 + 2}, \text{ for } n = 1, 2, \dots$$

Now

1.  $b_n \geq 0$ , for  $n = 1, 2, \dots$
2.  $b_n = \frac{n}{n^2 + 2} = \frac{1/n}{1 + 2/n^2} \rightarrow 0$  as  $n \rightarrow \infty$  so  $(b_n)$  is a null sequence
3.  $(b_n)$  is decreasing, because  $(1/b_n) = \left(\frac{n^2 + 2}{n}\right) = \left(n + \frac{2}{n}\right)$  is increasing.

Hence, by the Alternating Test,

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}n}{n^2 + 2} \text{ is convergent.}$$

## Solution to Additional Exercise D34

Since  $3n - 2 < 3n$  and  $3n - 1 < 3n$ , we have

$$\begin{aligned} \frac{1}{3n-2} + \frac{1}{3n-1} - \frac{1}{3n} &> \frac{1}{3n} + \frac{1}{3n} - \frac{1}{3n} \\ &= \frac{1}{3n}, \text{ for } n = 1, 2, \dots \end{aligned}$$

We now write series (1) as  $\sum_{n=1}^{\infty} a_n$  where

$$a_n = \frac{1}{3n-2} + \frac{1}{3n-1} - \frac{1}{3n}, \text{ for } n = 1, 2, \dots$$

We have just shown that

$$a_n > \frac{1}{3n}, \text{ for } n = 1, 2, \dots$$

Since  $\sum_{n=1}^{\infty} (1/n)$  is a basic divergent series, it follows from the Corollary to the Multiple Rule (Corollary D26) that  $\sum_{n=1}^{\infty} (1/(3n))$  is also divergent. It now follows from the Comparison Test that

$$\sum_{n=1}^{\infty} a_n \text{ is divergent.}$$

# Additional exercises for Unit D4

## Section 1

### Additional Exercise D35

Let  $f$  and  $g$  be the functions

$$f(x) = \tan x \quad (x \in (-\frac{1}{2}\pi, \frac{1}{2}\pi))$$

and

$$g(x) = \sqrt{x} \quad (x \in [0, \infty)).$$

Determine the domain and the rule of the following functions.

- (a)  $f + g$       (b)  $fg$       (c)  $f/g$       (d)  $f \circ g$   
 (e)  $g \circ f$

### Additional Exercise D36

Prove that the following function has an inverse function, and find the domain and rule of this inverse function.

$$f(x) = \frac{x-1}{x+2} \quad (x \in (-2, \infty))$$

*Hint:* It may help to write

$$\frac{x-1}{x+2} = 1 - \frac{3}{x+2}.$$

### Additional Exercise D37

By checking the definition, prove that each of the following functions is strictly monotonic.

- (a)  $f(x) = x^3 + 1 - \frac{1}{x^2} \quad (x \in (0, \infty))$   
 (b)  $f(x) = \frac{1}{(1+x^3)^2} \quad (x \in [0, \infty))$

## Section 2

### Additional Exercise D38

Use the appropriate rules, together with the list of basic continuous functions, to prove that the following functions are continuous. (Recall that  $\exp(x)$  is an alternative way of writing  $e^x$  and is useful when taking the exponential of a long expression.)

- (a)  $f(x) = \exp(\sin(x^2 + 1)) \quad (x \in \mathbb{R})$   
 (b)  $f(x) = e^{\sqrt{x}} + x^5 \quad (x \in [0, \infty))$

### Additional Exercise D39

Determine whether the following functions are continuous at 0.

- (a)  $f(x) = \begin{cases} x, & x < 0, \\ \frac{1}{x+1}, & x \geq 0. \end{cases}$   
 (b)  $f(x) = \begin{cases} \sin x \sin(1/x), & x \neq 0, \\ 0, & x = 0. \end{cases}$   
 (c)  $f(x) = \begin{cases} (1/x) \cos(1/x), & x \neq 0, \\ 0, & x = 0. \end{cases}$   
 (d)  $f(x) = \begin{cases} 0, & x < 0, \\ \sqrt{x}, & x \geq 0. \end{cases}$

### Additional Exercise D40

Prove that the following function is continuous at  $\frac{1}{2}\pi$  and at  $-\frac{1}{2}\pi$ .

$$f(x) = \begin{cases} -1, & x \leq -\frac{1}{2}\pi, \\ \sin x, & -\frac{1}{2}\pi < x < \frac{1}{2}\pi, \\ 1, & x \geq \frac{1}{2}\pi. \end{cases}$$

## Section 3

### Additional Exercise D41

Prove that the function

$$f(x) = x - \sin x - \frac{2}{3}\pi \quad (x \in \mathbb{R})$$

has a zero in the interval  $(\frac{2}{3}\pi, \frac{5}{6}\pi)$ .

### Additional Exercise D42

Suppose that the function  $f : [0, 1] \rightarrow [0, 1]$  is continuous. Prove that the equation

$$f(x) = x^3$$

has a solution in the interval  $[0, 1]$ .

*Hint:* Consider the function

$$g(x) = f(x) - x^3 \quad (x \in [0, 1]).$$

### Additional Exercise D43

Prove that each of the following polynomials has the stated number of zeros:

(a)  $p(x) = x^4 - 4x^3 + 3x^2 + 2x - 1$ , four zeros

(b)  $p(x) = 3x^3 - 8x^2 + x + 3$ , three zeros.

## Section 4

### Additional Exercise D44

Prove that each of the following functions has a continuous inverse function and determine the domain of the inverse function in each case.

(a)  $f(x) = x^3 + 1 - \frac{1}{x^2} \quad (x \in (0, \infty))$

(b)  $f(x) = \frac{1}{(1+x^3)^2} \quad (x \in [0, \infty))$

*Hint:* We proved that these functions are monotonic in Additional Exercise D37.

### Additional Exercise D45

State whether or not each of the following statements is true.

(a)  $\sin(\sin^{-1} x) = x$ , for  $x \in [-1, 1]$ .

(b)  $\sin^{-1}(\sin x) = x$ , for  $x \in \mathbb{R}$ .

### Additional Exercise D46

(a) Prove that the identity

$$\tan^{-1} x + \tan^{-1} y = \tan^{-1} \left( \frac{x+y}{1-xy} \right),$$

for  $x, y \in \mathbb{R}$ ,

is true, provided that  $\tan^{-1} x + \tan^{-1} y$  lies in  $(-\frac{1}{2}\pi, \frac{1}{2}\pi)$ .

(b) Hence evaluate

$$\tan^{-1}(\frac{1}{2}) + \tan^{-1}(\frac{1}{3}).$$

# Solutions to additional exercises for Unit D4

## Solution to Additional Exercise D35

(a) The domain of  $f + g$  is

$$(-\frac{1}{2}\pi, \frac{1}{2}\pi) \cap [0, \infty) = [0, \frac{1}{2}\pi);$$

the rule is

$$(f + g)(x) = \tan x + \sqrt{x}.$$

(b) As in part (a), the domain of  $fg$  is  $[0, \frac{1}{2}\pi)$ ; the rule is

$$(fg)(x) = \sqrt{x} \tan x.$$

(c) The domain of  $f/g$  is

$$[0, \frac{1}{2}\pi) - \{x : \sqrt{x} = 0\} = (0, \frac{1}{2}\pi);$$

the rule is

$$(f/g)(x) = \frac{\tan x}{\sqrt{x}}.$$

(d) The domain of  $f \circ g$  is

$$\{x \in [0, \infty) : \sqrt{x} \in (-\frac{1}{2}\pi, \frac{1}{2}\pi)\} = [0, \frac{1}{4}\pi^2);$$

the rule is

$$(f \circ g)(x) = \tan \sqrt{x}.$$

(e) The domain of  $g \circ f$  is

$$\{x \in (-\frac{1}{2}\pi, \frac{1}{2}\pi) : \tan x \in [0, \infty)\} = [0, \frac{1}{2}\pi);$$

the rule is

$$(g \circ f)(x) = \sqrt{\tan x}.$$

## Solution to Additional Exercise D36

First we solve the equation

$$y = \frac{x-1}{x+2}$$

to obtain  $x$  in terms of  $y$ . We find that

$$y = \frac{x-1}{x+2} = 1 - \frac{3}{x+2} \iff x = -2 + \frac{3}{1-y}.$$

Thus  $f$  is one-to-one, so  $f$  has an inverse function with rule  $f^{-1}(y) = -2 + 3/(1-y)$ .

Now we find the image set of  $f$ , which is the domain of  $f^{-1}$ . For each  $x \in (-2, \infty)$ , we have  $3/(x+2) > 0$ , so  $y < 1$ . Hence  $f((-2, \infty)) \subseteq (-\infty, 1)$ .

Also, for each  $y \in (-\infty, 1)$ , we have  $1-y > 0$ , so

$$x = -2 + \frac{3}{1-y} \in (-2, \infty).$$

Thus  $f((-2, \infty)) \supseteq (-\infty, 1)$ , so

$$f((-2, \infty)) = (-\infty, 1).$$

Hence the domain of  $f^{-1}$  is  $(-\infty, 1)$ , so, adopting the usual practice of denoting the domain variable by  $x$ , we have

$$f^{-1}(x) = -2 + \frac{3}{1-x} \quad (x \in (-\infty, 1)).$$

## Solution to Additional Exercise D37

(a) If  $0 < x_1 < x_2$ , then  $x_1^3 < x_2^3$  and  $x_1^2 < x_2^2$  so that  $-1/x_1^2 < -1/x_2^2$ .

Hence

$$x_1^3 + 1 - \frac{1}{x_1^2} < x_2^3 + 1 - \frac{1}{x_2^2},$$

so  $f$  is strictly increasing on  $(0, \infty)$ .

(b) If  $0 \leq x_1 < x_2$ , then  $0 \leq x_1^3 < x_2^3$ , so

$$1 + x_1^3 < 1 + x_2^3.$$

Hence

$$\frac{1}{(1+x_1^3)^2} > \frac{1}{(1+x_2^3)^2},$$

so  $f$  is strictly decreasing on  $[0, \infty)$ .

## Solution to Additional Exercise D38

(a) Let  $g(x) = x^2 + 1$ ,  $h(x) = \sin x$  and  $k(x) = e^x$ . Then  $g$ ,  $h$  and  $k$  are basic continuous functions. Hence, by the Composition Rule,

$$f = k \circ h \circ g \text{ is continuous.}$$

(b) Let  $g(x) = \sqrt{x}$ ,  $h(x) = e^x$  and  $k(x) = x^5$ . Then  $g$ ,  $h$  and  $k$  are basic continuous functions. Hence, by the Composition Rule and the Sum Rule,

$$f = h \circ g + k \text{ is continuous.}$$

## Solution to Additional Exercise D39

(a) We prove that

$$f(x) = \begin{cases} x, & x < 0, \\ \frac{1}{x+1}, & x \geq 0, \end{cases}$$

is discontinuous at 0, using Strategy D14. We choose

$$x_n = -\frac{1}{n}, \quad n = 1, 2, \dots$$

Then  $x_n \rightarrow 0$  and

$$f(x_n) = x_n \rightarrow 0.$$

Since  $f(0) = 1/(0+1) = 1$ , we have

$$f(x_n) \not\rightarrow f(0),$$

which shows that  $f$  is discontinuous at 0.

(b) We prove that

$$f(x) = \begin{cases} \sin x \sin(1/x), & x \neq 0, \\ 0, & x = 0, \end{cases}$$

is continuous at 0, by using the Squeeze Rule.

Now

$$-1 \leq \sin(1/x) \leq 1, \quad \text{for } x \neq 0,$$

so

$$-\sin x \leq \sin x \sin(1/x) \leq \sin x, \quad \text{for } x \in (0, \pi),$$

since  $\sin x \geq 0$ , for  $x \in (0, \pi)$ .

Also,

$$-\sin x \geq \sin x \sin(1/x) \geq \sin x, \quad \text{for } x \in (-\pi, 0),$$

since  $\sin x \leq 0$ , for  $x \in (-\pi, 0)$ .

Hence

$$-|\sin x| \leq \sin x \sin(1/x) \leq |\sin x|, \quad \text{for } 0 < |x| < \pi.$$

Since  $f(0) = 0$ , we deduce that

$$-|\sin x| \leq f(x) \leq |\sin x|, \quad \text{for } x \in (-\pi, \pi).$$

Thus, if we take  $I = (-\pi, \pi)$ , and

$$g(x) = -|\sin x| \quad \text{and} \quad h(x) = |\sin x|,$$

then condition 1 of the Squeeze Rule is satisfied.

Next,  $f(0) = g(0) = h(0) = 0$ , so condition 2 of the Squeeze Rule is satisfied.

Finally, the functions  $g$  and  $h$  are continuous at 0, by the Composition Rule and Multiple Rule, so condition 3 of the Squeeze Rule is satisfied.

Hence  $f$  is continuous at 0, by the Squeeze Rule.

(c) We prove that

$$f(x) = \begin{cases} (1/x) \cos(1/x), & x \neq 0, \\ 0, & x = 0, \end{cases}$$

is discontinuous at 0, using Strategy D14. We choose

$$x_n = \frac{1}{2n\pi}, \quad n = 1, 2, \dots$$

Then  $x_n \rightarrow 0$ , and

$$f(x_n) = \frac{1}{x_n} \cos\left(\frac{1}{x_n}\right) = 2n\pi \rightarrow \infty.$$

Since  $f(0) = 0$ , we have

$$f(x_n) \not\rightarrow f(0),$$

which shows that  $f$  is discontinuous at 0.

(d) We use the Glue Rule. Let  $I = \mathbb{R}$  and define the functions  $g$  and  $h$  by

$$g(x) = 0 \quad \text{and} \quad h(x) = \sqrt{x}.$$

Then  $f$  is defined on  $I$  and  $0 \in I$ . Also,

$$\begin{aligned} f(x) &= g(x), & \text{for } x \in (-\infty, 0), \\ f(x) &= h(x), & \text{for } x \in (0, \infty), \end{aligned}$$

so condition 1 of the Glue Rule holds with  $a = 0$ .

Moreover,  $f(0) = g(0) = h(0) = 0$ , so condition 2 holds.

Finally,  $g$  and  $h$  are basic continuous functions so they are continuous at 0, and so condition 3 holds.

Hence  $f$  is continuous at 0, by the Glue Rule.

## Solution to Additional Exercise D40

We use the Glue Rule. Take  $I = (-\frac{1}{2}\pi, \infty)$  and define the functions  $g$  and  $h$  by

$$g(x) = \sin x \quad \text{and} \quad h(x) = 1.$$

Then  $f$  is defined on  $I$  and  $\frac{1}{2}\pi \in I$ . Also,

$$\begin{aligned} f(x) &= g(x), & \text{for } x \in (-\frac{1}{2}\pi, \frac{1}{2}\pi), \\ f(x) &= h(x), & \text{for } x \in (\frac{1}{2}\pi, \infty), \end{aligned}$$

so condition 1 of the Glue Rule holds with  $a = \frac{1}{2}\pi$ .

Moreover,

$$g(\frac{1}{2}\pi) = f(\frac{1}{2}\pi) = h(\frac{1}{2}\pi) = 1,$$

so condition 2 holds.

Finally,  $g$  and  $h$  are basic continuous functions so they are continuous at  $\frac{1}{2}\pi$ , and so condition 3 holds.

Hence  $f$  is continuous at  $\frac{1}{2}\pi$ , by the Glue Rule.

A similar argument can be given for  $a = -\frac{1}{2}\pi$ , with  $I = (-\infty, \frac{1}{2}\pi)$ ,  $g(x) = -1$  and  $h(x) = \sin x$ .

(Alternatively, note that  $f$  is an odd function:

$$f(x) = -f(-x), \quad \text{for } x \in \mathbb{R},$$

so the continuity of  $f$  at  $-\frac{1}{2}\pi$  follows from its continuity at  $\frac{1}{2}\pi$ , by the Composition Rule and the Multiple Rule.)

### Solution to Additional Exercise D41

By calculation,

$$\begin{aligned} f\left(\frac{2}{3}\pi\right) &= \frac{2}{3}\pi - \sin\left(\frac{2}{3}\pi\right) - \frac{2}{3}\pi \\ &= -\sin\left(\frac{1}{3}\pi\right) = -\sqrt{3}/2 < 0 \end{aligned}$$

and

$$\begin{aligned} f\left(\frac{5}{6}\pi\right) &= \frac{5}{6}\pi - \sin\left(\frac{5}{6}\pi\right) - \frac{2}{3}\pi \\ &= \frac{1}{6}\pi - \sin\left(\frac{1}{6}\pi\right) = \frac{1}{6}\pi - \frac{1}{2} > 0, \end{aligned}$$

because  $\pi > 3$ .

Also,  $f$  is continuous by the Combination Rules (since it is a sum of basic continuous functions).

Hence, by the Intermediate Value Theorem,  $f$  has a zero in  $(\frac{2}{3}\pi, \frac{5}{6}\pi)$ .

### Solution to Additional Exercise D42

If  $f(0) = 0$  or  $f(1) = 1$ , then we can take  $c = 0$  or  $c = 1$ , respectively.

Otherwise, we have  $f(0) > 0$  and  $f(1) < 1$ , since  $0 \leq f(x) \leq 1$ , for  $0 \leq x \leq 1$ .

We consider the function

$$g(x) = f(x) - x^3 \quad (x \in [0, 1]).$$

Now  $g$  is continuous on  $[0, 1]$ , by the Combination Rules. Moreover

$$g(0) = f(0) - 0 > 0$$

and

$$g(1) = f(1) - 1 < 0.$$

Thus, by the Intermediate Value Theorem, there is a number  $c$  in  $(0, 1)$  such that

$$g(c) = 0, \quad \text{so} \quad f(c) = c^3.$$

### Solution to Additional Exercise D43

(a) When

$$p(x) = x^4 - 4x^3 + 3x^2 + 2x - 1,$$

we have

$$M = 1 + \max\{|-4|, |3|, |2|, |-1|\} = 5,$$

so all the zeros of  $p$  lie in  $(-5, 5)$ , by Theorem D54.

Calculating  $p(n)$  for integers  $n$  in  $[-5, 5]$ , we obtain

$n$	-2	-1	0	1	2	3
$p(n)$	55	5	-1	1	-1	5

Thus  $p$  changes sign on each of the intervals

$$[-1, 0], \quad [0, 1], \quad [1, 2], \quad [2, 3].$$

Since  $p$  is continuous, we deduce by the Intermediate Value Theorem that  $p$  has a zero in each of the intervals

$$(-1, 0), \quad (0, 1), \quad (1, 2), \quad (2, 3).$$

So  $p$  has at least four zeros. Since  $p$  is a polynomial of degree four, it has at most four zeros. Thus  $p$  has exactly four zeros.

(b) We have

$$\begin{aligned} p(x) &= 3x^3 - 8x^2 + x + 3 \\ &= 3\left(x^3 - \frac{8}{3}x^2 + \frac{1}{3}x + 1\right). \end{aligned}$$

For  $q(x) = x^3 - \frac{8}{3}x^2 + \frac{1}{3}x + 1$  we have

$$M = 1 + \max\left\{\left|-\frac{8}{3}\right|, \left|\frac{1}{3}\right|, |1|\right\} = \frac{11}{3},$$

so all the zeros of  $q$ , and hence of  $p$ , lie in  $(-\frac{11}{3}, \frac{11}{3})$ , by Theorem D54.

Calculating  $p(n)$  for integers  $n$  in  $[-4, 4]$ , we obtain

$n$	-2	-1	0	1	2	3
$p(n)$	-55	-9	3	-1	-3	15

Thus  $p$  changes sign on each of the intervals

$$[-1, 0], \quad [0, 1], \quad [2, 3].$$

Since  $p$  is continuous, we deduce by the Intermediate Value Theorem that  $p$  has a zero in each of the intervals

$$(-1, 0), \quad (0, 1), \quad (2, 3).$$

So  $p$  has at least three zeros. Since  $p$  is a polynomial of degree three, it has at most three zeros. Thus  $p$  has exactly three zeros.

### Solution to Additional Exercise D44

(a) We use Strategy D15.

1. We showed that  $f$  is strictly increasing on  $(0, \infty)$  in Additional Exercise D37(a).

2. The function

$$f(x) = x^3 + 1 - \frac{1}{x^2} = \frac{x^5 + x^2 - 1}{x^2} \quad (x \in (0, \infty))$$



is the restriction to  $(0, \infty)$  of a rational function which is continuous on  $\mathbb{R} - \{0\}$ . Hence  $f$  is continuous.

3. Choose the increasing sequence  $(n)$ , which tends to  $\infty$ , the right endpoint of  $(0, \infty)$ . Then

$$f(n) = n^3 + 1 - \frac{1}{n^2} \rightarrow \infty \text{ as } n \rightarrow \infty,$$

by the Reciprocal Rule. Thus the right endpoint of  $J = f((0, \infty))$  is  $\infty$ .

Then choose the decreasing sequence  $(1/n)$ , which tends to 0, the left endpoint of  $(0, \infty)$ . Then

$$f(1/n) = 1/n^3 + 1 - n^2 \rightarrow -\infty \text{ as } n \rightarrow \infty,$$

by the Reciprocal Rule. Thus the left endpoint of  $J$  is  $-\infty$ .

Hence  $J = (-\infty, \infty) = \mathbb{R}$ , so  $f$  has a continuous inverse function

$$f^{-1} : \mathbb{R} \longrightarrow (0, \infty),$$

by the Inverse Function Rule.

(b) We use Strategy D15.

1. We showed that  $f$  is strictly decreasing on  $[0, \infty)$  in Additional Exercise D37(b).
2. The function

$$f(x) = \frac{1}{(1+x^3)^2} \quad (x \in [0, \infty))$$

is the restriction to  $[0, \infty)$  of a rational function which is continuous on  $\mathbb{R} - \{1\}$ . Hence  $f$  is continuous.

3. We have  $f(0) = 1$ , so the corresponding endpoint of  $J = f([0, \infty))$  is 1, and  $1 \in J$ .  
Now choose the increasing sequence  $(n)$ , which tends to  $\infty$ , the right endpoint of  $[0, \infty)$ . Then

$$f(n) = \frac{1}{(1+n^3)^2} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Thus the corresponding endpoint of  $J$  is 0, and  $0 \notin J$ .

Hence  $J = (0, 1]$ , so  $f$  has a continuous inverse function

$$f^{-1} : (0, 1] \longrightarrow [0, \infty),$$

by the Inverse Function Rule.

## Solution to Additional Exercise D45

(a) True: by definition,

$$\sin(\sin^{-1} x) = x, \quad \text{for } x \in [-1, 1].$$

(b) False: for example,

$$\sin^{-1}(\sin 2\pi) = \sin^{-1} 0 = 0.$$

## Solution to Additional Exercise D46

(a) Let  $a = \tan^{-1} x$  and  $b = \tan^{-1} y$ . Then  $x = \tan a$  and  $y = \tan b$ , so

$$\tan(a+b) = \frac{\tan a + \tan b}{1 - \tan a \tan b} = \frac{x+y}{1-xy}.$$

Hence

$$a+b = \tan^{-1} \left( \frac{x+y}{1-xy} \right),$$

provided that  $a+b$  lies in  $(-\frac{1}{2}\pi, \frac{1}{2}\pi)$ , which is the image of  $\tan^{-1}$ . Thus

$$\tan^{-1} x + \tan^{-1} y = \tan^{-1} \left( \frac{x+y}{1-xy} \right),$$

provided that  $\tan^{-1} x + \tan^{-1} y$  lies in  $(-\frac{1}{2}\pi, \frac{1}{2}\pi)$ .

(b) Since  $\tan^{-1}$  is strictly increasing, we have

$$0 < \tan^{-1}(\frac{1}{2}) + \tan^{-1}(\frac{1}{3}) < 2 \tan^{-1}(1) = \frac{1}{2}\pi.$$

Hence, we can apply the formula in part (a):

$$\begin{aligned} \tan^{-1}(\frac{1}{2}) + \tan^{-1}(\frac{1}{3}) &= \tan^{-1} \left( \frac{\frac{1}{2} + \frac{1}{3}}{1 - \frac{1}{6}} \right) \\ &= \tan^{-1}(1) = \frac{1}{4}\pi. \end{aligned}$$